

Hyperacusis and Disorders of Sound Intolerance

Clinical and Research Perspectives

Editor-in-Chief for Audiology

Brad A. Stach, PhD

Hyperacusis and Disorders of Sound Intolerance

Clinical and Research Perspectives

Marc Fagelson, PhD
David M. Baguley, BSc, MSc, MBA, PhD





5521 Ruffin Road
San Diego, CA 92123

e-mail: info@pluralpublishing.com
website: <http://www.pluralpublishing.com>

Copyright © by Plural Publishing, Inc. 2018

Typeset in 11/13 Adobe Garamond by Achorn International
Printed in the United States of America by McNaughton & Gunn

All rights, including that of translation, reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, recording, or otherwise, including photocopying, recording, taping, Web distribution, or information storage and retrieval systems without the prior written consent of the publisher.

For permission to use material from this text, contact us by
Telephone: (866) 758-7251
Fax: (888) 758-7255
e-mail: permissions@pluralpublishing.com

Every attempt has been made to contact the copyright holders for material originally printed in another source. If any have been inadvertently overlooked, the publishers will gladly make the necessary arrangements at the first opportunity.

Library of Congress Cataloging-in-Publication Data

Names: Fagelson, Marc, editor. | Baguley, David (David M.), editor.
Title: Hyperacusis and disorders of sound intolerance : clinical and research perspectives / [edited by] Marc Fagelson, David M. Baguley.
Description: San Diego, CA : Plural Publishing, [2018] | Includes bibliographical references and index.
Identifiers: LCCN 2017058142 | ISBN 9781944883287 (alk. paper) | ISBN 1944883282 (alk. paper)
Subjects: | MESH: Hyperacusis—diagnosis | Hyperacusis—therapy | Tinnitus—diagnosis | Tinnitus—therapy
Classification: LCC RF293.7 | NLM WV 270 | DDC 617.8/9—dc23
LC record available at <https://lcn.loc.gov/2017058142>

Contents

Preface

vii

Contributors

ix

SECTION I. DEFINITIONS, MEASUREMENT, AND EPIDEMIOLOGY

- 1 Disorders of Sound Tolerance: History and Terminology 3
Marc Fagelson and David M. Baguley
- 2 Audiological Assessment of Decreased Sound Tolerance 15
Glynnis A. Tidball and Marc Fagelson
- 3 The Epidemiology and Natural History of Disorders of Loudness Perception 33
David M. Baguley
- 4 Scales and Questionnaires for Decreased Sound Tolerance 43
Kathryn Fackrell and Derek J. Hoare

SECTION II. MECHANISMS

- 5 Peripheral Mechanisms of Decreased Sound Tolerance 61
Roland Schaette
- 6 Tinnitus and Hyperacusis: Relationship, Mechanisms, and Initiating Conditions 77
Larry E. Roberts, Tanit Ganz Sanchez, and Ian C. Bruce
- 7 Hyperacusis: Medical Diagnoses and Associated Syndromes 107
Don McFerran
- 8 Animal Models of Hyperacusis and Decreased Sound Tolerance 133
Jos J. Eggermont
- 9 Traumatic Brain Injury and Auditory Processing 149
Melissa A. Papesh, Sarah M. Theodoroff, and Frederick J. Gallun
- 10 Psychological Aspects and Management of Hyperacusis 167
Gerhard Andersson

SECTION III. AUDITORY DISORDERS: MANIFESTATIONS

- 11 Reflections on the Association between Hyperacusis and Tinnitus 181
David M. Baguley

12	Diplacusis	191
	<i>Marc Fagelson</i>	

SECTION IV. MANAGEMENT

13	Increased Sound Sensitivity in Children	207
	<i>Veronica Kennedy, Claire Benton, and Rosie Kentish</i>	
14	Hearing Aids for Decreased Sound Tolerance and Minimal Hearing Loss: Gain Without Pain	223
	<i>Grant D. Searchfield and Caroline Selvaratnam</i>	
15	Hyperacusis Management: A Patient's Perspective	241
	<i>Rob Littwin</i>	
16	Hyperacusis: Past, Present, and Future	265
	<i>Marc Fagelson and David M. Baguley</i>	
	<i>Index</i>	271

Preface

For most of us, life is full of sound. While there are some individuals who lead a solitary, silent, or contemplative life, the more common path is replete with auditory stimulation. On an early morning walk in a North American city before conference presentations, the two editors of this book were intentionally attentive to the sounds around them, including sirens, engine sounds, phones ringing, and shouting. Many of the sound alerts in common use have been designed so as to be as compelling, and in a sense, as annoying as possible, cutting through distractions and other sounds to demand immediate and sustained attention (Patterson, 1990; Vastfjall et al., 2014).

For many years substantial scientific and clinical effort has been expended in understanding and ameliorating the impact of reduced hearing, resulting in sophisticated technologies such as hearing aids and auditory implants, advanced surgical techniques, and intensive rehabilitation strategies. Far, far less attention and scrutiny have been given to the experiences of individuals for whom the world of sound is more intense, more vivid, and perhaps perceived as more toxic than is usual. Most audiologists and otologists know of patients for whom everyday sound evokes discomfort, distress, aversion, and in some, pain.

This book seeks to explore and help the reader begin to understand those experiences. Some challenges will be encountered. The vocabulary used to describe such experiences is varied and imprecise, including decreased, reduced, or collapsed sound tolerance, and hyperacusis. Definitions of each of these terms varies; given such fundamental differences, it is not surprising that data regarding the epi-

demiology and natural history of hyperacusis is sparse, and inconsistent where it does exist. Some aspects of the physiological mechanisms of loudness or sound intensity perception in the auditory brain remain obscure, and there is a disconnect between the auditory neuroscience and the clinical communities that remains difficult to bridge. The experiences of individuals with reduced sound tolerance is heterogeneous, and can vary on a day to day, or hour to hour basis, and in some this is modulated by emotional and psychological state as well as the auditory environment. Tools to assess the extent and severity of loudness tolerance symptoms are crude, and in some cases may be deeply uncomfortable for the patient, as may be the case in some methods of ascertaining the threshold of loudness discomfort using sound stimulation. There is little in the way of hard evidence regarding therapy, and which interventions might be optimal for which type of patient.

All this may seem daunting, and lead one to consider that the topic of decreased sound tolerance cannot sensibly be addressed at all. However, looking at other fields would lead us to disagree. The field of pain studies was, until recently, in a similar state to that which is observed regarding hyperacusis (Chen, 2011), but systematic and focused endeavor has led to deeper understanding of mechanisms and the development of therapies that can be rigorously assessed in well-designed clinical trials. The present volume represents an attempt to take a first step on a similar journey, and it is our hope that it will inspire and provoke colleagues in both the auditory neuroscience and the clinical (especially audiology, otology,

and neurology) communities to discuss, explore, and research hyperacusis and related phenomena. Given the present state of knowledge, different authors have used different terminology, and have varied opinions, in their chapters: while we have edited with a goal of improving clarity and consistency, we have preserved some of these variations so that distinctive opinions, interpretations, and hypotheses might be heard.

Thanks are due to each of the authors for their fine contributions, and the time and effort expended on them, to Plural Publishing for agreeing to support this work, and to our families and colleagues for their support and forbearance. We would like to express our appreciation also to the many patients who have shared their experiences of hyperacusis and other phenomena of disordered loudness perception, from whom we have gleaned knowledge, experience, and inspiration.

—Marc Fagelson
David M. Baguley

References

- Chen, J. (2011). History of pain theories. *Neurosci. Bull.*, Oct, 27(5), 343–350. doi: 10.1007/s12264-011-0139-0.
- Patterson, R. (1990). Auditory warning sounds in the work environment. *Philos. Trans. R Soc. Lond. B Biol. Sci.*, 327(1241), 485–492.
- Vastfjall, D., Bergman, P., Sköld, A., Tajadura, A., & Larsson, P. (2012). Emotional responses to information and warning sounds. *J. Ergonomics*, 1, 106. doi:10.4172/2165-7556.1000106.

Note. DB is funded through the NIHR Biomedical Research Centre program, however, the views expressed are those of the authors and not necessarily those of the NIHR, the NHS, or the Department of Health.

Contributors

Gerhard Andersson, PhD, Dr. Med. Sci.

Professor in Clinical Psychology
Department of Behavioural Sciences
and Learning
Linköping University
Linköping, Sweden
Department of Clinical Neuroscience
Karolinska Institute
Stockholm, Sweden
Chapter 10

**David M. Baguley, BSc, MSc, MBA,
PhD, FAAA**

Professor of Hearing Services
President, British Tinnitus Association
Nottingham Biomedical Research Centre
University of Nottingham
Otology and Hearing Group
Division of Clinical Neuroscience
University of Nottingham
Honorary Consultant Clinical Scientist
(Audiology)
Nottingham University Hospitals NHS
Foundation Trust
Nottingham, United Kingdom
Chapters 1, 3, 11, and 16

Claire Benton, MSc

Consultant Clinical Scientist (Audiology)
Nottingham Audiology Services,
Nottingham University Hospitals NHS
Foundation Trust
Nottingham, UK
Chapter 13

Ian C. Bruce, PhD, PEng, FASA

Professor and Associate Chair for Graduate
Studies

Department of Electrical and Computer
Engineering
McMaster University
Hamilton, Ontario, Canada
Chapter 6

Jos J. Eggermont, PhD

Emeritus Professor
Physiology and Pharmacology, and
Psychology
University of Calgary
Calgary, Alberta, Canada
Chapter 8

Kathryn Fackrell, PhD

Research Fellow
University of Nottingham
Nottingham, United Kingdom
Chapter 4

Marc Fagelson, PhD

Professor
Department of Audiology and Speech
Language Pathology
East Tennessee State University
Johnson City, Tennessee
Chapters 1, 2, 12, and 16

Frederick J. Gallun, PhD

Research Investigator, VA RR&D
National Center for Rehabilitative Auditory
Research
Assistant Professor
Department of Otolaryngology—Head
and Neck Surgery
Oregon Health and Science University
Portland, Oregon
Chapter 9

Derek J. Hoare, PhD

Senior Research Fellow
University of Nottingham
Nottingham, United Kingdom
Chapter 4

**Veronica Kennedy, FRCS(Oto), FRCP,
FRCPC**

Consultant Audiovestibular Physician
Bolton NHS Foundation Trust
Bolton, United Kingdom
Chapter 13

Rosie Kentish, MA

Consultant Clinical Psychologist
Royal National Throat Nose and Ear
Hospital
London, United Kingdom
Chapter 13

Rob Littwin, MA

Musician
Producer
Arranger
Atlanta, Georgia
Chapter 15

Don McFerran, MA, FRCS

Consultant ENT Surgeon
Colchester Hospital
University NHS Foundation Trust
Colchester, United Kingdom
Chapter 7

Melissa A. Papesh, PhD, AuD

Research Investigator, VA RR&D
National Center for Rehabilitative Auditory
Research
Portland, Oregon
Chapter 9

Larry E. Roberts, PhD

Professor Emeritus
Department of Psychology Neuroscience
and Behaviour

McMaster University
Hamilton, Ontario, Canada
Chapter 6

Prof. Dr. Tanit Ganz Sanchez, MD, PhD

Associate Professor of Otolaryngology
University São Paulo School of Medicine
Founder of Instituto Ganz Sanchez
São Paulo, Brazil
Chapter 6

Roland Schaette, PhD

Lecturer
University College London, Ear Institute
London, United Kingdom
Chapter 5

**Grant D. Searchfield, BSc, MAud(Hons),
PhD**

Associate Professor, Audiology
Director, Hearing and Tinnitus Clinic
The University of Auckland
Member, The University of Auckland Centre
for Brain Research and Brain Research
New Zealand
Scientific Director, TinnitusTunes
Auckland, New Zealand
Chapter 14

**Caroline Selvaratnam, BSc, MAud,
MNZAS**

Audiologist
University of Auckland Clinics
Hearing and Tinnitus Clinic
University of Auckland
Auckland, New Zealand
Chapter 14

Sarah M. Theodoroff, PhD

Research Investigator, VA RR&D
National Center for Rehabilitative Auditory
Research
Assistant Professor
Department of Otolaryngology—Head
and Neck Surgery

Oregon Health and Science University
Portland, Oregon

Chapter 9

Glynnis A. Tidball, MSc, RAUD

Clinical Instructor

School of Audiology and Speech Sciences

University of British Columbia

Audiologist

Department of Audiology

St. Paul's Hospital

Vancouver, Canada

Chapter 2

Marc and David, as well as many other professionals and the contributors to this text, continue to learn what it means for people and their families to live with hyperacusis and other forms of sound intolerance; this book is dedicated to the many individuals, patients, and providers, who sparked and continue to nurture that interest.

SECTION I

Definitions, Measurement, and Epidemiology

1

Disorders of Sound Tolerance: History and Terminology

Marc Fagelson and David M. Baguley

Introduction

Traditionally the discipline of audiology has been concerned with the challenges faced by people with reduced hearing abilities in order to alleviate the effects associated with hearing loss. While there is a multitude of such people, and the burdens they face are certainly substantial, in recent years there has been increasing awareness of a population of people who have the experience of sound being too intense for them, rather than being too quiet. In this chapter we introduce this topic, and critically examine the various definitions and models that have been proposed, and identify where present knowledge is inadequate.

The impact of hearing loss, particularly of sensory and/or neural origin on the auditory system's loudness processing, is difficult to predict, and prone to the influence of many non-auditory factors. Although hearing aid use often improves an individual's communication ability and quality of life, the lack of a straightforward relation between a patient's sensitivity and their behavior on specific au-

diologic tests or routine activities precludes development of interventions for individuals with hearing impairment that are consistently as effective as, for example, glasses for myopia. A similar, complex and multifactorial relation appears to exist between disorders of loudness perception, their impact, and the handicap that is associated. This chapter will focus specifically on basic elements related to disorders of sound tolerance—items such as terminology and models, both auditory-based and psychology-based, that support the terminology—that continue to vex providers, patients, and students who seek clarity when managing patients in distress.

History

The concept that human hearing varies with the sound environment and context in which the person is listening has long been accepted, the English phrase, 'I could hear a pin drop' articulating the experience of intense and active listening. The experience of hearing becoming

acute at times of emotional intensity is also known, and used extensively in horror films: for example, the trope of increasing the loudness of an enemy's footsteps as they approach along a gravel drive is widespread. The fact that an increase in the perceived intensity of the sound environment can be involuntary is less well known, however. This forms a major plot device in the novel *The Woman in White* by Wilkie Collins (1860), although the symptom is un-named. The heroine of the novel is Miss Laura Fairlie, who as the novel opens has recently been orphaned, and has inherited some money, which renders her attractive to predatory suitors, and she seeks the protection of her uncle, Mr Frederick Fairlie. Unfortunately, he is unable to support her, as he is living as a reclusive invalid, due to a number of symptoms, one of which is a decreased ability to tolerate sound. In the first conversation between niece and uncle, held at some distance across a semi-darkened room, Frederick Fairlie articulates his situation thus:

Pray excuse me. But could you contrive to speak in a lower key? In the wretched state of my nerves, loud sound of any kind is indescribable torture to me. You will pardon an invalid?

Mr. Fairlie thus describes intense sound-evoked pain, leading to invalidity. While Laura eventually transcends the multiple adverse events that befall her, her uncle's situation, which is normally seen by literary critics as an abrogation of his male responsibilities, is a major setback.

Sound intolerance, such as that described by the unfortunate Mr. Fairlie, often accompanies hearing loss across patient groups of all ages, backgrounds, and auditory history, although it may also be present in individuals with hearing thresholds within the range of normal. In practice, this situation is described

in terms of dynamic range, or the signal intensities that comprise the range of values spanning the patient's threshold of sensitivity to their threshold of pain or discomfort. In hearing impaired listeners, the dynamic range is reduced in one of two ways. First, in the presence of loudness recruitment thresholds of sensitivity may be elevated while thresholds of discomfort remain essentially unchanged. Alternatively, in cases of hyperacusis, the threshold of discomfort may change over time, and become abnormally low; regardless of absolute sensitivity, the patient's dynamic range would be reduced. In extreme cases, perhaps such as that described by Mr. Fairlie, the patient's dynamic range may collapse and become limited to such a narrow range of intensities that the patient develops aversions to most routine sounds and curtails life activities accordingly.

No mention is made of Mr. Fairlie having a hearing loss, and he is probably an illustration of decreased sound tolerance with normal or age-appropriate hearing thresholds. In the case of a person with hearing loss, it is valuable to consider the paradox at work, as the presence of that hearing loss should, at least on the surface, allow a person to tolerate moderate sound levels without feeling discomfort. However, many patients with hearing loss experience inordinate loudness in the presence of what should be acceptable sound levels, powerful negative emotions in the presence of sounds that most other people do not find objectionable, or sounds that cause pain in and around the ears. Patients affected by such sounds may practice avoidance strategies to minimize their discomfort; we must also consider that strategy's shortcomings.

We hope to expand upon the idea, expressed in the book's preface, that patients would benefit from greater consensus among professionals regarding terms and definitions related to unusual sound experiences. If a patient avoids situations and environments

because of aversions to specific sounds or sound levels, then that patient's discomfort and emotional response could be exacerbated by seemingly inconsistent advice and counseling from health care providers. Accurate and effective counseling relies upon many factors; certainly, the list would include an agreed-upon lexicon. In this regard, audiologists have much to learn from the psychological and trauma literatures in which a variety of clinical approaches that target patients' beliefs and understanding take a primary role in the intervention. When basic terminology cannot be agreed upon, the potential for patient learning and adapting to a challenging condition likely decreases. In this chapter, we will review definitions and terms currently in use; our intent is to move the conversation forward in a way that prioritizes lexical items supporting the needs of patients and practitioners.

Hearing Loss and Loudness Growth

Loudness Recruitment

The development of the audiometer improved the precision with which auditory thresholds were estimated. Audiometers generated signals covering a broad range of frequency and intensity, and in addition to threshold measures facilitated identifying the relation between sound intensity and the perception of loudness. Early investigators (Fowler, 1936; Steinberg & Gardner, 1937) reported that patients with hearing loss experienced, in addition to elevated thresholds, an unusual relation between the intensity of a stimulus and the loudness it evoked. For some patients with hearing loss, a signal 10 dB SL regarding the puretone threshold (that

is, signal level 10 dB higher than the patient's threshold of hearing) was rated as a loudness (i.e., on a 1 to 100 scale) higher than the value provided by a normal-hearing person. Further, the rating provided by the listener with hearing loss increased by a greater amount than that of the normal hearing listener given identical increases in stimulus level. When loudness rating was graphed as a function of stimulus level, the hearing-impaired listener functions were steeper than those for normal hearing listeners. Investigators also found that, in general, at stimulus levels higher than 90 to 95 dB SPL, regardless of the hearing loss magnitude, the loudness functions as rated by hearing-impaired listeners approached and often could be superimposed upon those functions from normal hearing listeners. This loudness growth pattern was termed "loudness recruitment" by Fowler.

The loudness recruitment label captured a few interesting aspects of auditory system function. First, Fowler and other investigators employed the term to describe the way that an array of auditory nerve fibers would be influenced by damage to the cochlear hair cells. Given that not all fibers in the auditory nerve displayed identical thresholds, the loss of hair cells should have the most substantial effect on fibers with low thresholds, or those that would under normal conditions respond to audible signals of low intensity. The change in nerve innervation would commonly follow damage to some of the cochlea's most vulnerable components, the outer hair cells. While the sensory fiber population serving the outer hair cells would not be expected to influence audibility, the weakened contribution of the outer hair cells to basilar membrane mechanics would result in reduced stimulation of inner hair cells, thereby producing mild-moderate hearing loss.

At stimulus levels exceeding 50 to 60 dB, however, the relative contribution of the outer hair cell system to cochlear mechanics